


ApiLifeVar



INSTRUCTIONS FOR USE

WHAT DO I NEED TO CHECK BEFORE TREATING MY COLONIES WITH APILIFEVAR?

- Temperatures shall be between 64-95 degrees Fahrenheit.
- No suppers are installed.



KEY RULES FOR AN EFFECTIVE TREATMENT:

- During the autumn treatment, **treat as soon as possible after removing supers** (after the last honey harvest)
- **Treat all your colonies simultaneously** to avoid re-infestation.
- The recommended dosage is **3 tablets per colony** (need 1.5 sachets for a full treatment). **2 treatments per year can be made.**
- **Don't place the product in the center of the beehive or near the brood.** For best results treat colonies in the **late afternoon** or evening when the temperature is lower and the bees are in, or returning to the hive. The rate of sublimation of the active ingredients and the activity of the bee colony is lower and the bees will become accustomed to the odor more readily than if the product is applied at the hottest part of the day, when the bees are most active.

HOW TO USE APILIFEVAR?




1. Open the bag containing the two tablets.



2. Take one of the 2 tablets and break it into 4 pieces.

No need for extra space



3. Open the beehive and set the single pieces of the tablet around the edge of the brood nest on top of the top bars (in the corners of the beehive).

4. Repeat procedure as follow:

#1

1st application
Leave 7 to 10 days

#2

2nd application
Leave 7 to 10 days

#3

3rd application
Leave 12 days

You need 3 tablets (1.5 bags) to ensure a full treatment.

WARNINGS:

- The treatment used at temperatures over 95 degrees Fahrenheit may increase agitation and stress of bees. It is possible to observe an increase in the mortality of bees and bee brood.
- Bees tend to move in and take out the tablet from the hive or propolize the tablet. It is a normal behavior.
- Effectiveness is not guaranteed with a lower dose.



ALV-04-US-N01-03/17

Check out our other solutions to support honey bee health:

